

## CONCEPT QUICKSTART – Area under Simple Curves

**Unit:** Unit 8: Application of Integrals

**Subject:** For CBSE Class 12 Mathematics

---

### SECTION 1: UNDERSTANDING THE CONCEPT

In elementary geometry, we utilize static formulas to calculate the area of standard shapes like triangles, rectangles, and circles. However, as George David Birkhoff aptly noted (NCERT p. 292), "nature can be conceived in harmonious form" through Mathematics, and nature rarely presents itself in perfect polygons. To calculate the space enclosed by irregular, curved regions, we must transition from calculus-free geometry to Integral Calculus. Standard formulas are strategically inadequate here; integration allows us to solve for any region by viewing it as a dynamic accumulation of infinite parts. This shift is the definitive bridge between pure algebraic theory and the tangible physical world.

#### 1.1 What Is Area under Simple Curves?

The "Big Idea" is that a definite integral is not merely a numerical output of an algebraic process; it represents a geometric physical space. We utilize the **Strip Method**, where a region is visualized as being composed of a large number of infinitesimally thin vertical or horizontal strips. By expressing the area of one "elementary strip" as a differential ( $dA = y \, dx$ ), we use integration to sum these strips across a specific interval. Integration is essentially the "summation of strips," providing the exact area of curved regions that Euclidean geometry cannot define.

#### 1.2 Why It Matters

This topic serves as the primary real-world application of the integration techniques mastered in Chapter 7. It is the language used by engineers and physicists to calculate work done by variable forces, the center of mass of irregular components, and fluid pressure on curved surfaces. In the CBSE landscape, it serves as the ultimate test of a student's ability to combine graphical visualization with procedural rigor, proving that calculus is a practical tool for measuring the world around us.

#### 1.3 Prior Learning Connection

Success in this unit is built upon three strategic prerequisites:

- **Definite Integrals (Fundamental Theorem of Calculus):** Strategic success is impossible without the ability to evaluate  $F(b) - F(a)$ . This is the engine that converts your setup into a final numerical area.

- **Standard Curve Equations:** You must instantly recognize and sketch circles ( $x^2 + y^2 = a^2$ ), ellipses ( $x^2/a^2 + y^2/b^2 = 1$ ), and parabolas ( $y^2 = 4ax$ ). If you cannot sketch the boundary, you cannot set the limits.
- **Antiderivative Formulas:** Specifically, the "gatekeeper" formula for circles and ellipses:  $\int \sqrt{a^2 - x^2} dx$ . Mastery of this specific integral is the difference between a high score and a mid-range result.

#### 1.4 Core Definitions

- **[Elementary Area (dA)]**
- **NCERT Reference:** Chapter 8, Section 8.1, p. 293
- **Definition/Formula:**  $dA = y dx$  (for vertical strips) or  $dA = x dy$  (for horizontal strips)
- **Used In:** Foundational setup for all Problem Types (F1–F5).
- **[Area Bounded by a Curve (Vertical)]**
- **NCERT Reference:** Chapter 8, Section 8.2, p. 293
- **Definition/Formula:**  $A = \int_a^b f(x) dx$
- **Used In:** Basic Single Curve, Sign Change, Between Two Curves.
- **[Area Bounded by a Curve (Horizontal)]**
- **NCERT Reference:** Chapter 8, Section 8.2, p. 293
- **Definition/Formula:**  $A = \int_c^d g(y) dy$
- **Used In:** Mixed Boundary Regions, Symmetric Closed Curves (Alternative Method).
- **[Area of an Ellipse (Standard Form)]**
- **NCERT Reference:** Chapter 8, Section 8.2, p. 295
- **Definition/Formula:**  $y = (b/a)\sqrt{a^2 - x^2}$  for the first quadrant; Total Area =  $\pi ab$
- **Used In:** Symmetric Closed Curves.
- **[Absolute Value Principle]**
- **NCERT Reference:** Chapter 8, Section 8.2, p. 293 (Remark)
- **Definition/Formula:** Area =  $|\int_a^b f(x) dx|$  when  $f(x) < 0$
- **Used In:** Sign Change & Absolute Value problems where curves dip below the x-axis.

These definitions form the bedrock of the NCERT curricular framework, which we will now examine to understand the boundaries of your exam syllabus.

---

## SECTION 2: WHAT NCERT SAYS

The NCERT textbook acts as the definitive boundary for the CBSE syllabus. Mastering the specific examples and exercise logic within these pages is the most efficient path to exam success because the Board evaluates your ability to correctly translate these exact geometric descriptions into definite integrals.

### 2.1 Key Statements

1. **Vertical Strip Logic:** When the area is bounded by  $y = f(x)$  and the x-axis, we use vertical strips of height  $y$  and width  $dx$ .
2. **Horizontal Strip Logic:** If the curve is given as  $x = g(y)$  and bounded by the y-axis, horizontal strips (width  $dy$ ) are often simpler to calculate.
3. **Handling Negative Areas:** Since area is a physical quantity, if an integral result is negative (due to the curve being below the x-axis), the absolute value must be taken.
4. **Splitting Integrals:** If a curve crosses the x-axis at  $x = c$  within the bounds  $[a, b]$ , the total area is the sum of the absolute values of the integrals for  $[a, c]$  and  $[c, b]$ .
5. **Symmetry Advantage:** For standard circles and ellipses centered at the origin, you must calculate the area of one quadrant and multiply by four.

### 2.2 Examples and Exercises

- **Example 1 (Page 294): Area of a Circle**
  - **Strategic Objective:** Calculating  $x^2 + y^2 = a^2$  using symmetry and vertical strips.
  - **Impact:** A frequent examiner choice testing the "gatekeeper" formula:  $(x/2)\sqrt{(a^2 - x^2)} + (a^2/2)\sin^{-1}(x/a)$ .
  - **Exercise Mapping:** Exercise 8.1, Q3 (Difficulty: Easy).
- **Example 2 (Page 295): Area of an Ellipse**
  - **Strategic Objective:** Managing coefficients  $(b/a)$  in  $x^2/a^2 + y^2/b^2 = 1$ .
  - **Impact:** Tests algebraic discipline; forgetting the constant  $b/a$  is a common point of failure.
  - **Exercise Mapping:** Exercise 8.1, Q1, Q2 (Difficulty: Moderate).
- **Example 3 (Page 296): Line Crossing the Axis**
  - **Strategic Objective:** Calculating area for  $y = 3x + 2$  between  $x = -1$  and  $x = 1$ .
  - **Impact:** Tests the "zero-crossing" identification ( $x = -2/3$ ) and integral splitting.

- **Exercise Mapping:** Miscellaneous Exercise Q2, Q4 (Difficulty: Moderate to Hard).

Mastering the "what" of NCERT leads directly to the "how" of problem-solving, where pattern recognition becomes your most powerful tool.

---

### SECTION 3: PROBLEM-SOLVING AND MEMORY

Mathematics is essentially a game of "Pattern Recognition." By categorizing problems into specific "Types," you reduce exam-day anxiety and improve speed by providing yourself with a pre-defined mental roadmap for every question.

#### 3.1 Problem Types

- **Problem Type: Basic Single Curve (F1)**
- **Structural Goal:** Calculate the space under one function above the x-axis.
- **Recognition Cues:** "Area under the curve," "between ordinates  $x=a$  and  $x=b$ ,"  $f(x) \geq 0$ .
- **The "So What?" Layer:** You are simply finding the cumulative height of a moving vertical bar as it slides from  $a$  to  $b$ .
- **NCERT References:** Miscellaneous Exercise 1(i), 1(ii).
- **Confusable Types:** Contrast with "Sign Change" (F3); if the curve never goes negative, it is F1.
- **Problem Type: Symmetric Closed Curves (F2)**
- **Structural Goal:** Find the total area of a standard circle or ellipse.
- **Recognition Cues:** Equations like  $x^2 + y^2 = a^2$  or  $x^2/a^2 + y^2/b^2 = 1$ . No explicit bounds given.
- **The "So What?" Layer:** You calculate 25% of the shape (the first quadrant) and quadruple the result.
- **NCERT References:** Examples 1 & 2.
- **Confusable Types:** If only a "quadrant" is asked for (Exercise 8.1, Q3), do NOT multiply by four.
- **Problem Type: Sign Change (F3)**
- **Structural Goal:** Total area when the curve crosses the x-axis.
- **Recognition Cues:** Functions like  $\sin(x)$  or  $\cos(x)$  over a full period, or lines like  $y = 3x + 2$  where bounds include the crossing point.

- **The "So What?" Layer:** You treat "underground" areas as positive space by splitting the integral and using absolute value bars.
- **NCERT References:** Example 4 ( $\cos x$ ), Miscellaneous Exercise 3.
- **Confusable Types:** Basic Single Curve (where no crossing occurs).
- **Problem Type: Between Two Curves (F4)**
- **Structural Goal:** Calculate the area enclosed by two intersecting functions.
- **Recognition Cues:** "Area between  $y = f(x)$  and  $y = g(x)$ ," or "area enclosed by two parabolas."
- **The "So What?" Layer:** You are finding the height of a strip by subtracting the bottom curve from the top curve:  $[f(x) - g(x)]$ .
- **NCERT References:** Concept implied in Miscellaneous Exercise; essential for higher-order thinking.
- **Confusable Types:** Mixed Boundary Regions; if only one curve is present, it is F1 or F3.
- **Problem Type: Mixed Boundary Regions (F5)**
- **Structural Goal:** Areas bounded by a mix of lines, curves, and axes where horizontal strips are better.
- **Recognition Cues:** Curve given as  $x = g(y)$  or bounded by the  $y$ -axis (Exercise 8.1, Q4).
- **The "So What?" Layer:** You are switching your perspective to sum horizontal strips ( $dA = x dy$ ) from bottom to top.
- **NCERT References:** Exercise 8.1, Q4; Example 2 (Alternative Method).
- **Confusable Types:** Basic Single Curve (Vertical vs. Horizontal choice).

### 3.2 Step-by-Step Methods

#### Type: Basic Single Curve: Solution Method

- **Pre-Check:** Verify  $f(x)$  is continuous and  $f(x) \geq 0$  on the interval  $[a, b]$ .
- **Core Steps:**
  - **Step 1 (Setup):** Identify  $y = f(x)$  and bounds  $a, b$ . Sketch the region (Mandatory for marks!).
  - **Step 2 (Apply):** Formulate  $A = \int_a^b f(x) dx$ .
  - **Step 3 (Integrate):** Find the antiderivative  $F(x)$ .
  - **Step 4 (Conclude):** Evaluate  $F(b) - F(a)$ . State result in "square units."

- **Variants:** May involve polynomial, trigonometric, or exponential functions.
- **Boundary Conditions:** Do NOT use if the curve dips below the x-axis (Switch to F3 logic).

### Type: Symmetric Closed Curves: Solution Method

- **Pre-Check:** Confirm symmetry about both axes. Ensure the curve is centered at (0,0).
- **Core Steps:**
  - **Step 1 (Strategy):** State: "By symmetry, total area = 4 × Area in 1st Quadrant."
  - **Step 2 (Isolate):** Solve for y in terms of x (e.g.,  $y = \sqrt{a^2 - x^2}$ ).
  - **Step 3 (Integrate):** Set up  $4 \times \int_0^a f(x) dx$ .
  - **Step 4 (Scale):** Apply the limits and multiply by the 4x multiplier.
- **Variants:** Can be solved using horizontal strips ( $4 \times \int_0^b x dy$ ) with identical results.
- **Boundary Conditions:** If the curve is shifted (origin is not the center), the 4x multiplier may not apply without adjustment.

### 3.3 How to Write Answers

To earn full marks and avoid step-marking deductions, follow this standard **CBSE Answer Frame**:

- **Line 1 (Setup):** "The area bounded by the curve  $y = f(x)$ , the x-axis, and lines  $x = a$ ,  $x = b$  is  $A = \int_a^b f(x) dx$ ."
- **Line 2 (Identity):** " $A = \int_a^b$  [actual function]  $dx$ ."
- **Line 3 (Antiderivative):** " $= [F(x)]_a^b$ ."
- **Line 4 (Substitution):** " $= F(b) - F(a)$ ."
- **Line 5 (Result):** "[Final Value] square units."

**Essential Phrases:** "By symmetry about both axes," "The area element  $dA = y dx$ ," "Taking absolute value for the region below the x-axis."

#### ⚠ DEDUCTION WARNINGS:

- **Missing Sketch:** Loss of 0.5 to 1 mark.
- **Missing "Square Units":** Loss of 0.5 marks.
- **Forgetting b/a in Ellipse:** Loss of 1+ marks due to incorrect integration.

### 3.4 Common Mistakes (The "Pitfall" Gallery)

- **Pitfall 1: The Negative Area Trap**
  - **Category:** Logic.
  - **Symptom:** Reporting area as a negative number.
  - ✓ **Fix:** Area is physical space; always report the absolute value.
  - **Must-Check:** Always verify if the curve crosses the x-axis ( $f(x)=0$ ) between your bounds.
- **Pitfall 2: Limit Reversal**
  - **Category:** Algebra.
  - **Symptom:** Computing  $F(a) - F(b)$ .
  - ✓ **Fix:** Always Upper Limit minus Lower Limit:  $F(b) - F(a)$ .
- **Pitfall 3: The Multiplier Oversight**
  - **Category:** Logic.
  - **Symptom:** Finding the area of one quadrant but forgetting the 4x scale for the full circle/ellipse.
  - ✓ **Fix:** Clearly write "Total Area = 4 × Area in 1st Quadrant" as your very first step.
- **Pitfall 4: Top vs. Bottom (F4)**
  - **Category:** Logic.
  - **Symptom:** Subtracting the wrong curve, leading to negative results.
  - ✓ **Fix:** Always check the graph: Area =  $\int [f_{\text{top}}(x) - f_{\text{bottom}}(x)] dx$ .

### 3.5 Exam Strategy

1. **Foundational Phase:** Start with Exercise 8.1, Q1-Q3. These build speed with vertical strips.
2. **Symmetry Mastery:** Practice Examples 1 and 2 until the antiderivative for  $\sqrt{a^2 - x^2}$  is second nature. This is the "gatekeeper" formula for high-scoring circle/ellipse questions.
3. **Advanced Phase:** Move to the Miscellaneous Exercise to tackle trigonometric curves ( $\cos x$ ,  $\sin x$ ) and absolute value functions ( $|x|$ ). These require precise split-integral logic.

### 3.6 Topic Connections

This topic looks backward to **Chapter 7 (Integrals)** for its computational tools—you cannot find an area if you cannot integrate. It looks forward to **Physics and Engineering**, where these "areas" represent work, mass, and energy.

### 3.7 Revision Summary

- **Core Tool:** Integration is the "summation of strips."
- **Orientation:** Vertical ( $dx$ ) is standard; Horizontal ( $dy$ ) is a strategic choice for curves like  $x = g(y)$ .
- **Symmetry Rule:** Use  $4 \times \text{Area}(Q1)$  for standard circles and ellipses.
- **Positivity Rule:** Area must be non-negative; split integrals at zero-crossings.
- **Standard Formula:**  $\int \sqrt{a^2 - x^2} dx = (x/2)\sqrt{a^2 - x^2} + (a^2/2)\sin^{-1}(x/a)$ .
- **Setup Discipline Checklist:**
  1. Sketch the curve and shade the region.
  2. Identify the correct bounds ( $a, b$ ).
  3. Set up the integral form ( $\int y dx$  or  $\int x dy$ ).
  4. Check for sign changes (absolute value check).
  5. Integrate, substitute, and label with "square units."

Mastering the area under simple curves is not about complex integration, but about the disciplined setup of the elementary strip.

# Profsam.com